



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

XXXV. *Some New Properties in Conic Sections, discovered by Edward Waring, M. A. Lucasian Professor of the Mathematics in the University of Cambridge, and F R. S. to Charles Morton, M. D. Sec. R. S.*

T H E O R. I.

Read June 21,
1764.

SIT ellipsis APBQCRDSET, &c. describantur circa eam duo polygona [TAB. XIII. Fig. 1.] (*abcdef*, &c. *pqrstv*, &c.) eundem laterum numerum habentia, & quorum latera ad respectiva contactuum puncta (APBQCRDS, &c.) in duas æquales partes dividuntur, i. e. $aA = Ab$, $bB = Bc$, $cC = Cd$, &c. $pP = Pq$, $qQ = Qr$, $rR = Rs$, &c. & erit summa quadratorum ex singulis unius polygoni lateribus æqualis summæ quadratorum ex singulis alterius polygoni lateribus, i. e.

$$ab^2 + bc^2 + cd^2 + de^2 + ef^2 + \dots = p^2q^2 + qr^2 + rs^2 + st^2 + tv^2 + \dots$$

Cor. Ducantur lineæ AB, BC, CD, DE, EF, &c. PQ, QR, RS, ST, TV, &c. & erit $AB^2 + BC^2 + CD^2 + DE^2 + EF^2 + \dots = PQ^2 + QR^2 + RS^2 + ST^2 + TV^2 + \dots$

THEOR. II.

Iisdem positis fit O centrum ellipseos, & ducantur lineæ OA , OP , OB , OQ , OC , OR , OD , OS , &c. erit

$$OA^2 + OB^2 + OC^2 + OD^2 + \&c. = OP^2 + OQ^2 + OR^2 + OS^2 + \&c.$$

Cor. Ducantur etiam lineæ Oa , Op , Ob , Oq , Oc , Or , Od , Os , &c. & erit

$$Oa^2 + Ob^2 + Oc^2 + Od^2 + \&c. = Op^2 + Oq^2 + Or^2 + Os^2 + \&c.$$

Hæc etiam vera sunt de polygonis inter conjugatas hyperbolas eodem modo descriptis.

THEOR. III.

Sit conica sectio $MPQRSTM$ &c. [Fig. 2.] cujus diameter fit AL , et ejus ordinata ML ; fit $Mp = Mv$, & consequenter $Lp = Lv$.

Ducantur lineæ pq , qr , rs , st , tv , &c. quæ respective tangant conicam sectionem in punctis P , Q , R , S , T , &c. & erit contentum

$$pP \times qQ \times rR \times sS \times \&c. = Pq \times Qr \times Rs \times St \times Tv \times \&c. \text{ vel, quod idem est, summa omnium hujus generis rationum } (Pp : Pq, Qq : Qr, Rr : Rs, Ss : St, \&c.) \text{ erit nihilo æqualis.}$$

Cor. 1. Sit ellipsis $PQRSTV$ &c. circa eam describatur quodcunque polygonum ($pqrstuv$, &c.), [Fig.










Fig. 6.



[Fig. 3.] cujus latera respective tangant ellipſim in punctis P, Q, R, S, T, V, &c. & erit contentum

$$pP \times qQ \times rR \times sS \times tT \times vV \text{ \&c. } = Pq \times Qr \times Rs \times St \times Tv \times Vw \times \text{ \&c. }$$

Cor. Ducantur lineæ PQ, QR, RS, ST, &c. & pro finibus angulorum WP p , QP q , RQ r , QR s , SR s , TS t , &c. ſcribantur respective a, p, b, q, c, r, d, s , &c. & erit

$$abcd \text{ \&c. } = p q r s \text{ \&c. }$$

Et fic de polygonis inter conjugatis hyperbolas inſcriptis.

Idem verum eſt de polygono, cujus laterum ſumma vel area minima fit, circa quamcunque ovalem in ſeſe ſemper concavam deſcripto, ut conſtat e noſtra Miſcell. Anal.

T H E O R. IV.

Sit ellipſis PAQBRCSDTEVF, &c. [Fig. 4.] circa eam deſcribantur duo polygona $abcdef$, &c. $pqrstuv$, &c. eundem laterum numerum habentia; eorum latera ab, bc, cd, de, ef , &c. pq, qr, rs, st, tv , &c. respective tangant ellipſim in punctis A, B, C, D, E, F, &c. & P, Q, R, S, T, U, &c. & fit $aA : Ab :: pP : Pq$, & $bB : Bc :: qQ : Qr$ & $cC : Cd :: rR : Rs$ & $dD : De :: sS : St$, & fic deinceps. Et area polygoni $abcdef$, &c. æqualis erit areæ polygoni $pqrstuv$, &c.

Cor. Duo parallelogramma ($abcd$ & $pqrs$) circa datæ ellipſeos conjugatas diametros (AC & BD; PR, QS) [Fig. 5.] deſcripta, erunt inter ſe æqualia.

In hoc casu enim $aA = Ab$, $bB = Bc$, $cC = Cd$,
 $dD = Da$, & $pP = Pq$, $qQ = Qr$, $rR = Rs$,
 $sS = Sp$; & consequenter $aA : Ab :: pP : Pq$ &
 $bB : Bc :: qQ : Qr$, & sic deinceps: ergo per the-
 orema hæc duo parallelogramma erunt inter se æqualia,
 quæ est notissima ellipseos proprietas.

Idem dici potest de polygonis inter conjugatas hy-
 perbolas eodem modo descriptis.

T H E O R. V.

Rotetur conica sectio circa diametrum ejus (AL),

& sit $MA M'$, &c. solidum exinde generatum; sint
 pq , qr , rs , st , tv , vw , wp , &c. [Fig. 6.] lineæ,
 quæ tangant solidum in respectivis punctis P, Q, R, S,
 T, V, W, &c. & erit contentum

$$pP \times qQ \times rR \times sS \times tT \times vV \times wW \times \&c. = \\ Pq \times Qr \times Rs \times St \times Tv \times Vw \times \&c.$$

T H E O R. VI.

Sit ellipsis $APBQCR$, &c. rotetur circa diame-
 trum ejus BD; & circa conjugatas diametros (AC
 & BD, PR & QS) describantur elliptici cylindri
 ($pqrs$ & $acbd$) [Fig. 7.] solidum generatum cir-
 cumscribentes, & erunt hi duo cylindri inter se æ-
 quales.

Sint duo solida e truncatis conis composita, solidum
 generatum circumscribentibus, & quorum latera con-
 tinuo

tinuo eâdem ratione ad puncta contactuum dividuntur; erunt hæc duo solida inter se æqualia.

Et sic de solidis inter conjugatas hyperboloides eodem modo descriptis.

Facile constant plures consimiles conicarum sectionum proprietates.

Hujus generis proprietates affirmari possunt de infinitis aliis curvis, ut facile deduci potest e nostrâ Miscell. Anal.